

NADIS disease bulletins are written specifically for farmers, to increase awareness of prevalent conditions and promote disease prevention and control, in order to benefit animal health and welfare.
Farmers are advised to discuss their individual farm circumstances with their veterinary surgeon.

Rectal Prolapse

Compared to other farm species, the pig appears to be particularly vulnerable to prolapse of the rectal tissue through the anus which can be seen in any age group from as early as 1-2 days old up to adults. The fundamental cause of the prolapse is an increase in abdominal pressure, forcing a breakdown in the weak muscular support mechanism of the pelvis, which normally retains the rectum in place. There may well be both breed and gender differences in the vulnerability of individuals to prolapse occurring.

Causes of Prolapse

The following list provides an outline of the most common causes of prolapse, as seen in commercial pig farms:-

- 1) Diarrhoea or dysentery – particularly associated with large intestine inflammation that may include rectal inflammation (e.g. Salmonella, Swine Fever).
- 2) Constipation most likely to be seen in the adult close to farrowing.
- 3) Parturition – as a result of excessive straining.
- 4) Water shortage – leading to reduce water content of the faeces and increase straining to pass.
- 5) Medicines. Certain antibiotics (Tylosin, Lincocin) have been associated with oedema (swelling) of the lining of the rectum and subsequent prolapse. This is most likely seen with high doses.
- 6) Toxins. Some mycotoxins from feed or straw can be associated with rectal swelling and straining.
- 7) Rectal damage e.g. as a result of boars riding each other.
- 8) Coughing. The process of coughing causes an increase in abdominal pressure and, in some cases, this may be sufficient to push out the rectum. Many animals will expel faeces as they cough and the rectal lining will penetrate through the anus. In extreme cases, it does not return and remains prolapsed.
- 9) Fast growth. Prolapsing can often be a problem in fast growing pigs, particularly from 30-60kg on very high density diets.
- 10) Variable temperatures. Pigs have a poor ability to control their body temperature and tend to be adversely affected by variation in the ambient temperature and prone to chilling. The consequence is huddling and piling on top of one another. If a pig then coughs while another is lying on top of it, the abdominal pressure will be even higher than normal and the only place that the pressure can be relieved is at the anus.

Consequences

Once a prolapse has occurred, a number of events may follow:-

- 1) It rapidly returns into the anus.
- 2) It remains outside the anus and, due to the constrictive effect on blood and fluid drainage, it generally swells up. It is thus easily damaged by trauma on pen divisions, feeders etc.
- 3) It is eaten by other pigs in the pen. It is not uncommon to find blood in a pen and around the mouths of pigs but with no obvious prolapse in any other animals i.e. the prolapse will have been completely chewed off.

Long term consequences may be:-

- a) No effect – particularly if the prolapse returns without damage.
- b) Slow dying off of the prolapsed material over several weeks with the chances of secondary infection arising from rotting tissue.
- c) Rectal structure. The prolapse resolves but the scar tissue left forms a ring of slowly constricting tissue that eventually blocks the rectum leading to a “blown up” pig (Fig 2). Such animals require euthanasia.

Action

Any animal noticed with a prolapsed rectum should be isolated away from other pigs. If it is of slaughter weight, it can immediately be despatched for slaughter with a Schedule 18 casualty slaughter owners declaration. It should be transported in isolation.

If swollen but undamaged, it may be possible to replace the prolapse by sprinkling sugar or salt on it, leaving it 30 minutes and then gently pushing it back in. The osmotic effects of the salt/sugar draws out the fluid and shrinks the prolapse.

It may be necessary to place a purse string suture around the anus to retain the rectum once replaced.

In a large animal (sow), a rubber washing up glove placed over an undamaged prolapse may exert enough pressure to shrink the tissue and return it inside the rectum.

Where a prolapse is damaged and clearly not in a state to replace, it must be amputated. The easiest way to achieve this is to insert a pipe (1” diameter for a growing pig, 1 ½” for sows) into the prolapse and tie a ligature around the prolapse baring down onto the pipe. It is necessary to tie the pipe in with the loose ends of the ligature. This will cut the blood supply to the prolapsed material and allow it to dry up and drop off, usually in less than 7 days, although it may be necessary to re-tie the ligature after 3-4 days as the tissue shrinks. If corrugated pipe is available (e.g. electrical conduit), heavy duty rubber bands or even lamb elastrator rings can be used as a ligature.

In all cases, antibiotic cover should be provided.

Prevention

Clearly, prevention of rectal prolapses rests in being able to identify and correct the cause of the problem.

Copyright © NADIS 2004

Mark White BVSc DPM MRCVS