

Poultry(6) – Breeding & Egg Production

First Name:

Last Name:

Email Address:

Veterinary Practice:

Postcode:

Please circle one answer only e.g. A

1. Eggs for artificial incubation should be:

- A. warm
- B. dirty
- C. clean
- D. cracked

2. Egg laying is influenced by:

- A. the cockerel
- B. light levels
- C. light nestboxes
- D. feeding grain

3. Hatchability is improved by:

- A. breeder ration
- B. feeding wheat
- C. two cockerels
- D. heavy feathering

4. Downgraded eggs have:

- A. thick shells
- B. meat spots
- C. regular shape
- D. brown shells

5. Vent gleet is:

- A. common
- B. curable
- C. a venereal herpesvirus
- D. a respiratory disease

6. The pin bones indicate:

- A. a lazy bird
- B. a laying bird
- C. a different breed
- D. a cockerel

7. Poor hatchability is caused by:

- A. clean eggs
- B. cheap incubator
- C. natural hatching
- D. poor nutrition

8. Egg peritonitis is caused by

- A. poor egg production
- B. yolk in the abdomen
- C. white in the abdomen
- D. no cockerel

9. Fertility is reduced by

- A. heavy feathering
- B. breeder ration
- C. longer days
- D. breeding ratio of 1:4

10. A disease which affects eggs is

- A. Mycoplasma
- B. Campylobacter
- C. Vent gleet
- D. Infectious Bronchitis