

Skin Conditions in Cattle - Ectoparasites

First Name:

Last Name:

Email Address:

Veterinary Practice:

Postcode:

Please circle one answer only e.g. A

1. Louse populations are highest:

- A. in cattle kept indoors during the winter months
- B. in cattle kept outdoors during the winter months
- C. in cattle kept indoors during the summer months
- D. in cattle kept outdoors during the summer months

2. Spread of lice occurs by:

- A. direct contact with infested cattle
- B. contact with wildlife such as badgers
- C. direct contact with contaminated surfaces such as handling facilities
- D. direct contact with sheep

3. Treatment of all louse species and internal parasites can be achieved using a:

- A. Group I anthelmintic such as fenbendazole
- B. Group II anthelmintic such as levamisole
- C. Group III anthelmintic such as pour-on ivermectin
- D. topical application of high cis cypermethrin or deltamethrin

4. Infestation with *C. bovis* typically affects which area of the body?

- A. the base of the tail
- B. the neck
- C. the legs
- D. the head

5. Midges are important in the transmission of which disease?

- A. BVD
- B. Foot and mouth disease
- C. Bluetongue
- D. Salmonellosis

6. In certain parts of the UK and Ireland, ticks are important vectors for which of the following diseases?

- A. BVD
- B. Foot and mouth disease
- C. Redwater
- D. Salmonellosis